

Plasma sheet structure in the magnetotail: kinetic simulation and comparison with satellite observations

Paolo Ricci^(1,2), Giovanni Lapenta^(1,3) and J. U. Brackbill⁽⁴⁾

(1) Istituto Nazionale per la Fisica della Materia (INFN), Dipartimento di Fisica, Politecnico di Torino, Torino, Italy

(2) Dipartimento di Energetica, Politecnico di Torino, Torino, Italy

(3) Theoretical Division, Los Alamos National Laboratory, Los Alamos NM, USA

(4) Department of Mathematics and Statistics, University of New Mexico, Albuquerque NM

P. Ricci, Dipartimento di Energetica, Politecnico di Torino, Corso Duca degli Abruzzi 24 - 10129 Torino, Italy.

G. Lapenta, Los Alamos National Laboratory, Los Alamos NM 85744 (lapenta@lanl.gov).

J.U. Brackbill, Department of Mathematics and Statistics, University of New Mexico, Albuquerque NM.

Abstract.

We use the results of a three-dimensional kinetic simulation of an Harris current sheet to shown and reproduce the ISEE-1/2, Geotail, and Cluster observations of the magnetotail current sheet structure. Current sheet flapping, current density bifurcation, and reconnection are explained as the results of the self-consistent evolution of a Harris current sheet, where lower-hybrid drift, kink, and tearing instabilities are involved.

1. Introduction

The magnetotail current sheet is one of the key topics in magnetospheric physics.

Observations of the current sheet have revealed a very complex structure. At the end of April 2, 1978, the ISEE-1/2 spacecraft detected a flapping of the plasma sheet and the spacecraft crossed the central region more than 10 times in a hour. In particular, during a "turbulent" crossing, the spacecraft detected current concentration outside the central region, unlike the Harris current sheet [Sergeev *et al.*, 1993]. In fact, Geotail [Kokobun *et al.*, 1994; Mukai *et al.*, 1994] averaged data obtained from October 1993 to June 1995 show that the structure of the plasma sheet can be often approximated by a double-peaked electric current sheet [Hoshino *et al.*, 1996] and observations made by the same spacecraft during a substorm on 23 April 1996 lead to a similar conclusion [Asano *et al.*, 2003]. On January 14, 1994, Geotail also detected multiple double-peaked current sheet crossings, associated with plasma flow [Hoshino *et al.*, 1996]. More recently, time analysis of data from the four Cluster spacecrafts [Balogh *et al.*, 2001] showed that fast motion and bifurcation of the current sheet are associated with a wave-like transient propagating in the dawn-to-dusk direction [Sergeev *et al.*, 2003; Runov *et al.*, 2003]. Plasma flow has also been observed during a substorm event [Hoshino *et al.*, 1996; Øieroset *et al.*, 2001; Asano *et al.*, 2003]. These observations refer both to the distant magnetotail ($\approx 100R_E$) [Hoshino *et al.*, 1996] and to a region closer to Earth ($\approx 15R_E$) [Sergeev *et al.*, 1993; Asano *et al.*, 2003; Runov *et al.*, 2003; Sergeev *et al.*, 2003]

A useful but simple one-dimensional description of the current sheet is given by the Harris model, where the magnetic field is given by $B_x(z) = B_0 \tanh(z/\lambda)$ and the plasma

density, proportional to the current density, is given by $n(z) = n_0 \cosh^{-2}(z/\lambda)$, where λ is the half thickness of the current sheet and the GSM coordinates are used.

Generalizations of the standard Harris current sheet equilibrium have been recently proposed to reproduce the bifurcation observed by satellites [*Shindler and Birn*, 2002; *Sitnov et al.*, 2003]. *Runov et al.* [2003] propose that bifurcation and flapping signatures are due to a kink or a sausage instability of a current sheet with an enhanced current density on both hemispheres. *Zelenyi et al.* [2002] show that non-adiabatic effects can reduce the current density in the center of the current sheet. A bifurcated current sheet can be present in the plasma outflow region when magnetic reconnection is occurring [e.g., *Arzner and Sholer*, 2001]. *Karimabadi et al.* [2003a, 2003b] argue that the ion-ion kink instability causes a displacement of the current sheet which can explain the flapping observations and interpret the bifurcated structure of the current sheet as a magnetic field profile with weak central gradient due to the non-linear evolution of the kink instability. Generally, plasma flow is explained in terms of plasma out-flowing from a reconnection region.

The aim of the present work is to analyze the results of a three-dimensional kinetic simulation of the Harris current sheet by introducing diagnostic tools very similar to the one used by satellite. We show that the self-consistent evolution of the current sheet displayed by the simulations can be responsible for the data observed by the satellites described in the references above. In particular, taking into account the relative motion of the current sheet and the spacecraft, and the Cluster tetrahedron configuration, we recover the most significant magnetic data records obtained by the Cluster spacecraft as

the signature of current sheet flapping. The recurrence frequency of the magnetic field B_x allow the comparison with observations by GEOTAIL [*Hoshino et al.*, 1996] that show current bifurcation, and the magnetic field structure show the signature of bifurcation recovered in single crossing. We also compare the plasma flow due to the tearing instability with the observations.

2. The simulations

In our study, we use the implicit PIC code CELESTE3D [*Forsslund and Brackbill*, 1985; *Vu and Brackbill*, 1992; *Ricci et al.*, 2002a], which is particularly suitable for large scale and long period kinetic simulations performed with high mass ratio and has been applied previously to problems in magnetospheric physics [e.g., *Lapenta and Brackbill*, 2000; *Lapenta and Brackbill*, 2002; *Ricci et al.*, 2002b; *Lapenta et al.*, 2003].

We use the same plasma parameters as the GEM challenge [*Birn et al.*, 2001]. In particular we start from a standard Harris current sheet, with magnetic field given by $B_x(z) = B_0 \tanh(z/\lambda)$, density by $n(z) = n_0 \cosh^{-2}(z/\lambda) + n_b$ with $\lambda = 0.5c/\omega_{pi}$, $T_i/T_e = 5$, the ion drift velocity is $V_{i0} = 1.67V_A$, and a background population $n_b = 0.2n_0$. Unlike the GEM challenge, we do not add any initial perturbation and let the system evolve on its own. The dimensions of the system are $[-L_x/2, L_x/2] \times [-L_y/2, L_y/2] \times [-L_z/2, L_z/2]$ with $L_x = 12.8c/\omega_{pi}$, $L_y = 19.2c/\omega_{pi}$, and $L_z = 6.4c/\omega_{pi}$, discretized with a grid $N_x \times N_y \times N_z = 32 \times 48 \times 32$. The boundary conditions are perfect conductors at $z = \pm L_z$ and periodic boundaries in all the other directions. The mass ratio is $m_i/m_e = 180$. The parameters chosen make the current sheet particularly unstable and its dynamics are accelerated compared with typical magnetotail current sheets. We are constrained to do that in

order to follow the dynamics of the current sheet in a reasonable computational time. As a consequence, it is necessary to scale our results to make a quantitative comparison between simulation results and observations. In any case, the general trends can be located and the linear theory [Karimabadi *et al.*, 2003a] can help in scaling the results.

Previous simulations [Lapenta and Brackbill, 2002; Lapenta *et al.*, 2003; Daughton, 2003] performed in the current aligned plane show, in absence of a plasma background, the development of the fastest lower-hybrid drift instability on the electron gyroscale, followed by electromagnetic modes with wavelengths intermediate between the ion and the electron gyroscale. The lower hybrid drift instability causes a velocity shear (present since the beginning of the simulation when a background plasma is present) that triggers a Kelvin-Helmholtz (KH) instability that kinks the current sheet. As we add a background population, following Karimabadi *et al.* [2003b], the velocity shear is present since the beginning of the simulation and the resulting KH instability can be also interpreted as a kinetic ion-ion kink instability [Karimabadi *et al.*, 2003a; Karimabadi *et al.*, 2003b].

Both the present and previous simulations have shown that with the kinking of the current sheet a tearing instability grows [Lapenta and Brackbill, 2001; Lapenta *et al.*, 2003]. In the present case, the fastest short wavelength modes grow (corresponding to $k_x L \approx 0.5$ or $m_x = 2$ in our simulation box), and then they merge to form a single island tearing mode that involve the whole domain.

Below we consider specific aspects of the satellite observations basing their interpretation on the simulation.

3. Current sheet flapping

When vertical oscillations of the plasma sheet (flapping) occur, spacecrafts may repeatedly cross the current sheet. Clear evidence of current sheet flapping is shown by ISEE-1/2 [Sergeev *et al.*, 1996], by Geotail [Hoshino *et al.*, 1996], and by Cluster [Runov *et al.*, 2003; Sergeev *et al.*, 2003]. We focus our attention on the observations by Cluster and show that current sheet kinking developed in the course of our simulations can explain those.

Figure 1 shows fully developed current sheet kinking. The B_x field is shown. The wavelength is $k_y\lambda \approx 0.5$, which matches fairly well the observed wavelength in Runov *et al.* [2003] ($k_y\lambda = 0.7$). The linear theory predicts a decrease of the wavelength when ρ_i/λ increases [Karimabadi *et al.*, 2003a] consistent with the fact that our thickness is likely smaller than the observation. The amplitude A at time $t\omega_{ci} = 16$ is $A/\lambda \approx 2$ is comparable to the observed value ($A/\lambda \approx 1.4$) [Sergeev *et al.*, 2003]. The flapping motion observed by Cluster moving duskward at $v_{ph} \approx 200$ km/s, corresponding to approximately $0.2v_A$. The kink instability shown in our simulations gives a $v_{ph,SIM} \approx 0.5v_A$, larger than observed in space. However, the linear theory predicts a decrease of the phase velocity when ρ_i/λ increases. Since we use an artificially high ρ_i/λ , the higher phase speed is justified and consistent with our interpretation of the flapping motion.

In Fig. 2a we show Cluster #2 and #3 observations taken on 29 August 2001, which have been analyzed previously by Runov *et al.* [2003]. In particular, the B_x data is considered. In Fig. 3a, we evaluate the magnetic field as a function of time as would be recorded by a virtual spacecraft placed in the environment provided by the simulation. According to the real spacecraft disposition, we impose a distance between the two virtual

satellites in the z direction to be of the order of $\lambda/2$. Cluster observes an oscillation period of $\tau = 90\text{s}$ and a relative velocity between satellite and plasma $v_{ph} \approx 0.2v_A$. In order to decrease the time necessary for the observation, we increase the relative satellite velocity up to $v_{SIM} = 5v_A$, thus decreasing the oscillation period to $\tau_{SIM} = 2\omega_{ci}^{-1}$, in good agreement with the oscillation period recorded by Cluster. With the new relative velocity and using the fact that $\omega_{ci} \approx 0.6\text{s}^{-1}$ in the magnetotail and the observed period is of the order $54\omega_{ci}^{-1}$, the observed wavelength, $v_{ph}\tau \approx 11c/\omega_{pi}$ and the simulated wavelength, $v_{SIM}\tau_{SIM} \approx 10c/\omega_{pi}$, are comparable.

The flapping observation recorded by Cluster #3 on September 26, 2001 and described by Sergeev *et al.* [2003] is shown in Fig. 2b. It is reproduced by our simulations at later times, when the amplitude of the kink has grown enough that the virtual satellite can pass from one side to the other of the current sheet. This is shown in Fig. 3b. We note that Cluster observations reveal a flattening of the current sheet in the vicinity of the points where $B_x = 0$, which is associated to current sheet bifurcation. The grid spacing used in the three-dimensional simulation does not allow to resolve this structure, which is better addressed in more resolved two-dimensional simulations.

In agreement with Sergeev *et al.* [2003] and Runov *et al.* [2003], our simulations reveal that the current sheet flapping is mostly in the (y, z) plane, while the tilt in the (x, z) plane is insignificant.

4. Current sheet bifurcation

Experimental evidence exists that the current distribution in the magnetotail sheet may be double-peaked, with a pair of electric current sheets separated by a layer of a weak

quasi-uniform magnetic field. Current sheet bifurcation has been revealed both in averages over a number of current sheet crossings, and in single sheet crossings.

The statistical studies of the current sheet presented in Hoshino *et al.* [1996] reveal a bifurcated current profile. An ensemble of neutral sheet crossings is considered and the occurrence frequency of B_x is evaluated. The observed distribution has a peak around the null magnetic region, as in Sergeev *et al.* [2003]. The distribution of occurrence of the field B_x , $N(B_x)$ can be expressed as

$$N(B_x) \propto \frac{d}{dB_x} F^{-1}(B_x) \quad (1)$$

where $B_x(z) = F(z)$. Thus, the functional form of the magnetic field as a function of z can be obtained, and from the gradient of B_x with respect to z it is possible to evaluate the plasma current. The whole procedure is able to smear out current sheet flapping and particular motion of the current sheet.

In order to study current bifurcation, we have performed a two-dimensional simulation in the (y, z) plane that allows to use a more refined grid in this plane ($N_y \times N_z = 128 \times 64$). In Fig. 4a we show the plot of in-plane current $\sqrt{J_y^2 + J_z^2}$ at $t\omega_{ci} = 20$. Although the high fluctuations, the simulation show an increase of the current on the flanks of the current sheet. To follow GEOTAIL data analysis, we compute the recurrence frequency of B_x (Fig 4b). The occurrence frequency shows a peak around $B_x = 0$, as satellite data show [Hoshino *et al.*, 1996 (Fig.2), Sergeev *et al.*, 2003 (Fig. 4)]. By integrating Eq. (1), we find $B_x(z)$ (Fig. 4c), which is compared with a Harris sheet profile, and the current profile as a function of z (Fig. 4d). The current is depleted at the center of the current sheet

and two current peaks grow on the flanks of the initial current sheet. (This is unlike the Harris sheet equilibrium, where $\partial B_x/\partial z$ is maximum at $z = 0$ where $B_x = 0$.) In Fig. 4d the current density profile is compared with GEOTAIL observations [*Hoshino et al.*, 1996 (Fig.4)] and found in remarkable agreement.

Single crossing observations of current sheet bifurcation are shown by Runov *et al.* [2003] and by Sergeev *et al.* [2003]. We focus on Fig. 3c in Sergeev *et al.* [2003], which shows reduced $\partial B_x/\partial z$ in the central part of the current sheet (reduced current) and enhanced gradient (enhanced current density) at the boundary part. In Fig. 5, we plot a number of B_x profiles as a function of z , at different values of y . The same features of the magnetic field structure shown by satellite observations is recovered.

We finally remark that observations by Geotail on 23 April 1996 show that positive $d|B_x|/dt$ corresponds to relevant current density J_y [*Asano et al.*, 2003]; the same effect is also recovered within our simulation.

5. Reconnection

Not only does a kink instability grow, but also a tearing instability develops in the Harris sheet, which leads to the reconnection of the magnetic field lines and outflow and inflow plasma jets. Satellite observations typically reveal the reconnection process either by detecting inflow and outflow plasma jets, which can be very noisy [e.g., *Asano et al.*, 2003], or by detecting earthward and tailward plasma jets with velocities of the order of $0.1v_A$ or bigger [*Hoshino et al.*, 1996], or even by detecting flow reversal [*Øieroset et al.*, 2001]. In Fig. 6 we show signatures of magnetic reconnection by showing a flow reversal associated with a change in the sign of the reconnecting field. The earthward and

tailward velocities, detected during the crossing of the current sheet, are of the order of $0.1v_A$, which roughly corresponds to the typical order of magnitude of plasma velocity in the satellite observations.

6. Conclusion

We have used the results of three-dimensional and two-dimensional kinetic simulations of Harris current sheet to show that satellite observations of current sheet flapping, current bifurcation, and reconnection can all be explained self-consistently. We have chosen to start from a relatively thin and unstable current sheet ($\lambda/d_i = 0.5$) in order to accelerate the plasma dynamics. Such thin current sheets are indeed observed in the magnetotail [e.g., *Asano et al.*, 2003, for a review].

Flapping oscillations have been shown as the results a kink instability that affects the whole current sheet dynamics. Frequency and amplitude compare well with satellite observations. Both average and single crossing signatures of current sheet bifurcation have been detected. Flow reversal, signature of reconnection, is also shown in presence of a changing sign B_z component.

Acknowledgments. The authors gratefully thank M. Hoshino for the permission to use the data plotted in Fig. 4 and J. Birn, J. Chen, W. Daughton, I. Furno, M. Taylor, A. Vaivads for helpful discussions. The satellite data has been obtained from Cluster FGM team [*Balogh et al.*, 2001]. This research is supported by the Laboratory Directed Research and Development (LDRD) program at the Los Alamos National Laboratory, by the United States Department of Energy, under Contract No. W-7405-ENG-36 and by NASA, under the "Sun Earth Connection Theory Program".

References

- Arzner, K., and M. Sholer, Kinetic structure of the post plasmoid plasma sheet during magnetic reconnection, *J. Geophys. Res.*, *106*, 3827, 2001.
- Asano, Y., et al., Evolution of the thin current sheet in a substorm observed by Geotail, *J. Geophys. Res.*, *108*(A5), 1189, doi: 10.1029/2002JA009785, 2003.
- Balogh, A., et al., The Cluster magnetic field investigation: Overview of in-flight performance and initial results, *Ann. Geophys.*, *19*, 1207, 2001.
- Birn, J. et al., Geospace Environment Modelling (GEM) magnetic reconnection challenge, *J. Geophys. Res.*, *106*, 3715, 2001.
- Brackbill, J. U. and D. W. Forslund, Simulation of low frequency, electromagnetic phenomena in plasmas, in *Multiple time Scales*, J.U. Brackbill and B.I. Cohen Eds., (Accademic Press, Orlando, 1985), pp. 271-310.
- Daughton, W.S., Electromagnetic properties of the lower-hybrid drift instability in a thin current sheet, *Phys. Plasmas*, submitted, 2003.
- Hoshino, M., et al., Structure of plasma sheet in magnetotail: Double-peaked electric current sheet, *J. Geophys. Res.*, *101*, 24775, 1996.
- Lapenta, G., and J.U. Brackbill, 3D reconnection due to oblique modes: a simulation of Harris current sheets, *Nonlinear Processes Geophys.*, *7*, 151, 2000.
- Lapenta, G., and J.U. Brackbill, Nonlinear evolution of the lower hybrid drift instability: Current sheet thinning and kinking, *Phys. Plasmas*, *9*, 1544, 2002.
- Lapenta, G., J.U. Brackbill, and W.S. Daughton, The unexpected role of the lower hybrid drift instability in magnetic reconnection in three dimensions, *Phys. Plasmas*, *10*, 1577,

2003.

Karimabadi, H., P.L. Pritchett, W. Daughton, and D. Krauss-Varban, Ion-ion kink instability in the magnetotail 1. Linear Theory, *J. Geophys. Res.*, in press (2003a).

Karimabadi, H., P.L. Pritchett, W. Daughton, and D. Krauss-Varban, Ion-ion kink instability in the magnetotail: 2. Three-dimensional full particle and hybrid simulations and comparison with observations, *J. Geophys. Res.*, in press (2003b).

Kokobun, S., et al., The GEOTAIL magnetic field experiment, *J. Geomagn. Geoelectr.*, *46*, 4, 1994.

Mukai, T.S., et al., The low energy particle (LEP) experiment on board the GEOTAIL satellite, *J. Geomagn. Geoelectr.*, *46*, 669, 1994.

Øieroset, M, T.D. Phan, M. Fujimoto, R.P. Lin, R.P. Lepping, In situ detection of collisionless reconnection in the Earth's magnetotail, *Nature*, *412*, 414, 2001.

Ricci, P., G. Lapenta, and J.U. Brackbill, A simplified implicit Maxwell solver, *J. Comput. Phys.*, *183*, 117, 2002a.

Ricci, P., G. Lapenta, and J.U. Brackbill, GEM reconnection challenge: Implicit kinetic simulations with the physical mass ratio, *Geophys. Res. Lett.*, *29*(23), 2088, doi:10.1029/2002GL015314, 2002b.

Runov, A., et al, Cluster observation of a bifurcated current sheet, *Geophys. Res. Lett.*, *30*(2), 1036, doi:10.1029/2002GL016136, 2003.

Sergeev, V.A., D.G. Mitchell, C.T. Russel, and D.J. Williams, Structure of the Tail Plasma/Current sheet at $\approx 11 R_E$ and its changes in the course of a substorm, *J. Geophys. Res.*, *98*, 17345, 1993.

- Sergeev, V., et al., Current sheet flapping motion and structure observed by Cluster, *Geophys. Res. Lett.*, *30*(6), 1327, doi: 10.1029/2002GL016500, 2003.
- Shindler, K., and J. Birn, Models of two-dimensional embedded thin current sheets from Vlasov theory, *J. Geophys. Res.*, *107*(A8), doi: 10.1029/2001JA000304, 2002.
- Sitnov, M.I., P.N. Guzdar, and M. Swisdak, A model of the bifurcated current sheet, *Geophys. Res. Lett.*, *30*(13), 1712, doi: 10.1029/2003GL017218, 2003.
- Vu, H. X. and J. U. Brackbill, CELEST1D: An implicit, fully kinetic model for low-frequency Electromagnetic plasma simulation, *Comput. Phys. Commun.*, *69*, 253, 1992.
- Zelenyi, L.M., D.C. Delcourt, H.V. Malova, and A.S. Sharma, "Aging" of the Magnetotail thin current sheet, *Geophys. Res. Lett.*, *29*(12), doi:10.1029/2001GL013789, 2002.

- Fig. 1: The kink of the current sheet is presented by showing the x component of magnetic field, B_x . Both quantities are shown as a function of y and z , at time $t\omega_{ci} = 16$ and at $x = 0$. B_x is normalized to B_0 .

- Fig. 2: Signatures of current sheet flapping, observed by the FGM Cluster experiment [Balogh *et al.*, 2003]. We report the B_x magnetic field recorded by satellites #2 (dashed) and #3 (solid) on 29 August 2001 that has been described by Runov *et al.* [2003] (a), and by satellite #3 on September 26 2001, described by Sergeev *et al.* [2003] (b).

- Fig. 3: Signatures of current sheet flapping as would be recorded by a virtual spacecraft placed in the environment provided by the simulation and which reproduce the real signature shown in Fig. 2. The B_x magnetic field is plotted, normalized to B_0 .



- Fig. 4: Current density $\sqrt{J_y^2 + J_z^2}$ from the two-dimensional simulation at time $t\omega_{ci} = 20$ (a), B_x (normalized to B_0) occurrence frequency (b), B_x profile as a function of z (solid) and comparison with Harris current sheet (dotted) (the normalization is arbitrary) (c), and current profile from the simulation compared with Geotail observations [Hoshino *et al.*, 2003, Fig. 4b] (the original dimensionless units have been scaled to fit the simulation results) (d).


- Fig. 5: B_x profile as a function of z for different value of y .

- Fig. 6: Typical signature of reconnection: during the crossing of the current sheet, the reconnecting field, B_z , changes sign (a) and it is associated to earthward and inward plasma jets (b).

This figure "fig1.JPG" is available in "JPG" format from:


<http://arXiv.org/ps/physics/0308034v3>






This figure "fig4.JPG" is available in "JPG" format from:

<http://arXiv.org/ps/physics/0308034v3>



a)



b)

